

Lumiprobe Corporation

201 International Circle, Suite 135 Hunt Valley, Maryland 21030

USA

Phone: +1 888 973 6353 Fax: +1 888 973 6354 Email: order@lumiprobe.com

BDP® 558/568 DBCO


http://www.lumiprobe.com/p/bdp-558-568-dbco

BDP 558/568 fluorescence maximum is in the yellow spectrum range (569 nm). This dye is an analog of such fluorophore as $Cy3^{\text{TM}}$ by its spectral characteristics. It is soluble in polar organic solvents and has good photostability and high quantum yield.


BDP 558/568 has hydrophobic properties and can be used for staining lipids, membranes, and other lipophilic compounds.

Introduced in the BDP 558/568 molecule, the cyclooctyne functional group allows entering into reactions of 1,3-dipolar cycloaddition with various substituted azides. Such reactions are promoted by ring strain and do not require copper catalysis. Thus, this method can be used for investigating various processes in living cells.

BDP 558/568 DBCO can be used for the detection of target molecules, proteins, or nucleic acids that contain azide groups by microscopy and flow cytometry.



Structure of BDP 558/568 DBCO



Absorption and emission spectra of BDP 558/568

General properties

Appearance: brown powder with luster

Molecular weight: 646.56

Molecular formula: $C_{37}H_{33}N_4BF_2O_2S$

Solubility: very soluble in DMF, DMSO, dichloromethane

Quality control: NMR ¹H, HPLC-MS (95%)

Storage conditions: Storage: 24 months after receival at -20°C in the dark. Transportation: at room

temperature for up to 3 weeks. Avoid prolonged exposure to light.

Legal statement: This Product is offered and sold for research purposes only. It has not been tested for

safety and efficacy in food, drug, medical device, cosmetic, commercial or any other use. Supply does not express or imply authorization to use for any other purpose, including, without limitation, in vitro diagnostic purposes, in the manufacture of food

or pharmaceutical products, in medical devices or in cosmetic products.

Spectral properties

Excitation/absorption maximum, nm: 561 ϵ , L·mol⁻¹·cm⁻¹: 84400 Emission maximum, nm: 569 Fluorescence quantum yield: 0.68